

85% -SUCCESS RATE 25 fields 81 wells 166 well tests

4D field gravity & 1D well log gravity observations

3D joint inversion of field and well log gravity (4D+1D)

3D model of density with
1 m vertical resolution

 3D model of current gas saturation after
10 years of production

3D model of current formation pressure after 10 years of production



PRODUCTION MONITORING

OVER TWO DECADES OF RELIABLE HYDROCARBON-BEARING RESERVOIR EXPLORATION

MAPPING PRESSURE DROP AND FLOODING AREAS BASED ON THE RESULTS OF TIME-LAPSE GRAVIMETRIC OBSERVATIONS AND BOREHOLE GRAVIMETRY (4D+1D)

Nadym-Pur, Western Siberia, Russian Federation Time lapse gravity survey: 1998-2008

GEOLOGICAL PROBLEM

- Monitoring of waterflooding in the main gas pool Cenomanian, Cretaceous, $K_{1\text{-}2} pk$
- Seismic monitoring not an option due to the absence of a seismic signal above the structure
- Study of more shallow reservoir in Berezov Formation, Senonian, Upper Cretaceous, K₂br

3D GRAVITY INVERSION WORKFLOW

Structural framework of the model was built using 2D & 3D seismic data interpretation, including target Cenomanian and Senonian horizons of Upper Cretaceous. Wells were used to define the initial 3D density model of target Cenomanian intervals (Upper Cretaceous, K₁). 3D density model was refined by joint 3D linear inversion of surface gravity field and well log gravity data with one-meter resolution in depth (Figure 1). We used surface background measurements of 1998 and measurements after a decade of production of 2008. 3D model of gas saturation (Figure 3) was built using the dependency of Cenomanian rock density on porosity and saturation (Figure 5). The dependency of real gas density on pressure and temperature was derived (Figure 4) as well as the dependency of Cenomanian gassaturated sandstone's density on porosity and gas saturation (Figure 5). Gas saturation decrease was detected in the marginal part of the gas pool which is associated with contour water flood within highly permeable beds (Figure 3). In the apical part of the structure, there is a zone of abnormal pressure drop (Figure 6). New gas-saturated pool was mapped in Cenonian formation of Upper Cretaceous (Figure 2).

GEOLOGICAL RESULTS

- Local areas of pressure drop by 5-6 MPa and gas-saturation change after 10 years of production were identified
- Predicted location of well clusters with gas-water contact rise and pressure drop
- Predicted gas-saturated reservoir in Senonian formation



Figure 6. Pressure change in Cenomanian gas pool after 10 years of production



Figure 1. High-resolution (1m in depth) 3D density models of gas-saturated reservoir by joint gravity field and gravity log inversion: background model (1998, left) and model after 10 years of production (2008, right)



Figure 2. Cross-section of 3D density model through the wells with gravity log



Figure 3. Cross-section of 3D gas-saturation model through the wells with gravity log



Figure 4. Dependency of gas density on preassure and temperature in Cenomanian gas-saturated sandstones



Figure 5. Dependency of Cenomanian rock density on porosity, water- and gas-saturation